

Measurement of Reactive Nitrogen in the Troposphere.

J. Sen, G. B. Osterman, G. C. Toon, R. J. Salawitch, J. J. Blavier,
J. J. Margitan, D. B. Peterson H. M. Pickett, R. A. Stachnik (Jet
Propulsion Laboratory, MS 183-301, California Institute of
Technology, 4800 Oak Grove Drive, Pasadena, CA 91109; 818-
354-1165; e-mail: sen@mark4sum.jpl.nasa.gov)

G. K. Yue (NASA Langley Research Center, MS 475, Hampton,
VA 23681)

High resolution infrared spectra made in September 1993 at 35°
N with the JPL MkIV interferometer have been used to deter-
mine the volume mixing ratio of nitrogen oxide gases (NO , NO_2 ,
 HNO_3 , HNO_4 , N_2O_5 , and ClONO_2) in the troposphere and lower
stratosphere (5 - 20 km). We present estimates of the N oxidation
ratio, HNO_3/NO_x , as well as profiles for HNO_4 , which constrains
our understanding of nitrogen and hydrogen radicals. Measured
concentrations of reactive nitrogen gases will be compared to
theoretical values computed using a photochemical model con-
strained by simultaneous MkIV observations of O_3 , NO_y , H_2O ,
 CH_4 , Cl_y , CO , C_2H_6 , and aerosol surface area from SAGE II.

1996 Fall Meeting

2. 008874470
- 3 (a) J. Sen,
Jet Propulsion Laboratory,
M. S. 183-301,
California Inst. of Tech.,
4800 Oak Grove Drive,
Pasadena, CA 91109-8099
(b) Tel: (818)354-1165
(b) Fax: (818)354-5148
- 4 A
- 5 (a) A02
- Tropospheric CO, NO_y & O_3
(b) 0355
- Troposphere
- composition and chemistry
6. Oral Preferred
- 7.
- 8 00% G (submitted)
9. Citeque enclosed
10. C
- 1 1
- 12.
- 3 3